

Updated on: 24th March 2023

CERTIFICATE OF ANALYSIS

Lot#: CyHum17030-HE-C

PRODUCT DESCRIPTION

Reference: HuHECSM/4-Product: Cryopreserved Human Hepatocytes

Category: Suspension Spheroid qualified: NO

(see details below: 3D Spheroid formation section)

Isolation date: 15th December 2017 Initial Isolation Viability: 81% Storage conditions: -196°C using LN₂

Sterility test: negative for bacteria, yeast, and

fungi

DONOR DEMOGRAPHICS

| Species | Gender | Race | Age | ВМІ | Smoker | Alcohol Use | Drug Use | | |
|------------------|-----------|-----------|---|-------------------------------|--------|-------------|----------|--|--|
| Human | Male | Caucasian | 49 | 23.39 | No | No | No | | |
| Р | Pathology | | | Serological Data ¹ | | | | | |
| Metastatic tumor | | | Tested negative less than 3 months before surgery | | | | | | |


Patient informed consent was obtained. ¹The donor was serologically tested negative for following infectious diseases: HIV, Hepatitis B and C. Donor medical history was also examined prior to accepting this donor. For donor's medication information, please contact us.

CHARACTERIZATION FOR SUSPENSION CELLS

| Post Thaw Lot information | Result | SD | n |
|------------------------------|----------------------|------------------------|---|
| Number of viable cells/vial: | 3.66x10 ⁶ | ± 1.16x10 ⁶ | 4 |
| Viability (%): | 86.72 | ± 2.61 | 4 |

Human hepatocytes were thawed according to Cytes Biotechnologies protocol. The post-thawing yield and viability (trypan blue exclusion assay) of hepatocytes were assessed after a purification process.

| Time (h) | 0 | 0.5 | 1 | 1.5 | 2 | 3 | 4 | 5 |
|---------------|--------|--------|--------|--------|--------|--------|--------|--------|
| Viability (%) | 85.10 | 83.15 | 83.47 | 85.64 | 82.15 | 82.23 | 78.27 | 74.26 |
| SD | ± 0.87 | ± 3.62 | ± 0.90 | ± 2.02 | ± 5.83 | ± 1.91 | ± 0.30 | ± 6.72 |



Resuspended human hepatocytes suspension (0,5 * 10⁶ cells in 0.5 ml medium) from the post-thaw assessment were incubated up to 5 h at 37°C. The assay was performed in 2 ml round-bottom tubes under shaking conditions (1000 rpm) using Eppendorf Thermomixer C. In the first two hours, samples were taken at every 30 min, after 2 h samples were taken at every 60 min. At each time point the viability of cells was calculated.

For basic research use only, not to be used for clinical or diagnostic applications. Products distributed by Cytes Biotechnologies may contain human material that should be treated as potentially biohazardous.



3D SPHEROID FORMATION

Spheroid morphology

Cytes **does not guarantee** that these primary hepatocytes will be suitable for 3D culture and creation of spheroid structures.

If you need help for an experiment, just contact us, our experts will be pleased to assist you

CERTIFICATION:

The viability and performance of the primary human hepatocytes provided depend primarily on the use of appropriate media and reagents, as well as the use of sterile plastics. Likewise, proper handling protocols must be followed. Please note that if these parameters are not carefully considered, the cellular response obtained in the assays may be lower than expected.

| Name | Tittle | Signature | Cytes Biotechnologies, S.L. | Date |
|---------------------------|-----------------|-------------|------------------------------|----------|
| Pilar Sainz de la Maza | Quality Manager | Par Jan Len | CYTES BOTECHPOLOGIES S.L. | 24/03/23 |

For basic research use only, not to be used for clinical or diagnostic applications. Products distributed by Cytes Biotechnologies may contain human material that should be treated as potentially biohazardous.